

## Atrial Fibrillation

### **What is atrial fibrillation?**

Atrial fibrillation, often called AFib or AF, is the most common type of treated heart arrhythmia. An arrhythmia is when the heart beats too slowly, too fast, or in an irregular way.

When a person has AFib, the normal beating in the upper chambers of the heart (the two atria) is irregular, and blood doesn't flow as well as it should from the atria to the lower chambers of the heart (the two ventricles). AFib may happen in brief episodes, or it may be a permanent condition.

Atrial Fibrillation is the most common type of heart arrhythmia. AFib occurs when the upper chambers and lower chambers are not coordinated, causing the heart to beat too slowly, too quickly, or irregularly.

### **Facts About AFib**

- It is estimated that 12.1 million people in the United States will have AFib in 2030.
- In 2018, AFib was mentioned on 175,326 death certificates and was the underlying cause of death in 25,845 of those deaths.
- People of European descent are more likely to have AFib than African Americans.
- Because the number of AFib cases increases with age and women generally live longer than men, more women than men experience AFib.

### **What are the symptoms of AFib?**

Some people who have AFib don't know they have it and don't have any symptoms. Others may experience one or more of the following symptoms:

- Irregular heartbeat
- Heart palpitations (rapid, fluttering, or pounding)
- Lightheadedness
- Extreme fatigue
- Shortness of breath
- Chest pain

### **What are the risk factors for AFib?**

The risk for AFib increases with age. High blood pressure, the risk for which also increases with advancing age, accounts for about 1 in 5 cases of AFib.

Risk factors for AFib include

- Advancing age
- High blood pressure
- Obesity
- European ancestry
- Diabetes
- Heart failure
- Ischemic heart disease
- Hyperthyroidism
- Chronic kidney disease
- Moderate to heavy alcohol use
- Smoking
- Enlargement of the chambers on the left side of the heart

### **How is AFib related to stroke?**

AFib increases a person's risk for stroke. When standard stroke risk factors were accounted for, AFib was associated with a four- to fivefold increased risk of ischemic stroke. AFib causes about 1 in 7 strokes.

Strokes caused by complications from AFib tend to be more severe than strokes with other underlying causes. Strokes happen when blood flow to the brain is blocked by a blood clot or by fatty deposits called plaque in the blood vessel lining.

### **How is AFib treated?**

Treatment for AFib can include

- Medicines to control the heart's rhythm and rate
- Blood-thinning medicine to prevent blood clots from forming and reduce stroke risk
- Surgery
- Medicine and healthy lifestyle changes to manage AFib risk factors

Article Source

Centers for Disease Control (CDC)

Source URL

<https://www.cdc.gov>

Last Reviewed

Thursday, October 22, 2020